



PCT/EP200 4 / 0 0 4 2 4 5



INVESTOR IN PEOPLE

10/563803

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

REC'D 03 JUN 2004

WIPO

PCT


The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



R. Mahoney

Signed

Dated 18 March 2004



Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office
Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference

MG/JW/PB60227P

2. Patent application number

(The Patent Office will fill in his part)

0309222.8

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

Glaxo Group Limited
Glaxo Wellcome House, Berkeley Avenue,
Greenford, Middlesex UB6 0NN, Great Britain

473587003
United Kingdom

4. Title of the invention

Novel Compounds

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent
(including the postcode)

Patents ADP number (if you know it)

Corporate Intellectual Property

GlaxoSmithKline
Corporate Intellectual Property (CN9 25.1)
980 Great West Road
BRENTFORD
Middlesex TW8 9GS

8072555006

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or each of these earlier applications and (if you know it) the or each application number

Country	Priority application number (if you know it)	Date of filing (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application	Date of filing (day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer yes if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is named as an applicant, or
- c) any named applicant is a corporate body

See note (d)

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form
Description
Claim(s)
Abstract
Drawings

15 ✓
2 ✓

C₁

10. If you are also filing any of the following, state how many against each item.

Priority Documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents
(*please specify*)

11.

We request the grant of a patent on the basis of this application

Signature

M Gibson

Date 23-Apr-03

12. Name and daytime telephone number of person to contact in the United Kingdom

M Gibson 01279 644841

Warning

After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission unless an application has been filed at least six weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505
- b) Write your answers in capital letters using black ink or you may type them.
- c) If there is not enough space for all relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- d) If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- f) For details of the fee and ways to pay please contact the Patent Office.

Patents Form 1/77


NOVEL COMPOUNDS

The present invention relates to novel piperidine carbonyl piperazine derivatives having pharmacological activity, processes for their preparation, to compositions containing them and to their use in the treatment of neurological and psychiatric disorders.

WO 97/06802 (Zeneca Limited) describe a series of pyridyl and pyrimidyl derivatives as oxido-squalene cyclase inhibitors which are claimed to lower blood cholesterol.
WO 02/76925 (Eli Lilly), WO 03/004480, WO 03/024928 and WO 03/024929 (all Novo Nordisk A/S and Boehringer Ingelheim International) describe a series of substituted piperidines which are claimed to bind selectively to the histamine H3 receptor.

The histamine H3 receptor is predominantly expressed in the mammalian central nervous system (CNS), with minimal expression in peripheral tissues except on some sympathetic nerves (Leurs *et al.*, (1998), Trends Pharmacol. Sci. **19**, 177-183). Activation of H3 receptors by selective agonists or histamine results in the inhibition of neurotransmitter release from a variety of different nerve populations, including histaminergic and cholinergic neurons (Schlicker *et al.*, (1994), Fundam. Clin. Pharmacol. **8**, 128-137). Additionally, *in vitro* and *in vivo* studies have shown that H3 antagonists can facilitate neurotransmitter release in brain areas such as the cerebral cortex and hippocampus, relevant to cognition (Onodera *et al.*, (1998), In: The Histamine H3 receptor, ed Leurs and Timmerman, pp255-267, Elsevier Science B.V.). Moreover, a number of reports in the literature have demonstrated the cognitive enhancing properties of H3 antagonists (e.g. thioperamide, clobenpropit, ciproxifan and GT-2331) in rodent models including the five choice task, object recognition, elevated plus maze, acquisition of novel task and passive avoidance (Giovanni *et al.*, (1999), Behav. Brain Res. **104**, 147-155). These data suggest that novel H3 antagonists such as the current series could be useful for the treatment of cognitive impairments in diseases such as Alzheimer's disease and related neurodegenerative disorders.

The present invention provides, in a first aspect, a compound of formula (I) or a pharmaceutically acceptable salt thereof:



wherein:

- R¹ represents aryl, heteroaryl, -aryl-X-aryl, -aryl-X-heteroaryl, -aryl-X-heterocyclyl, -heteroaryl-X-heteroaryl, -heteroaryl-X-aryl or -heteroaryl-X-heterocyclyl; wherein said aryl, heteroaryl and heterocyclyl groups of R¹ may be optionally substituted by one or more (eg. 1, 2 or 3) substituents which may be the same or different, and which are selected from the group consisting of halogen, hydroxy, cyano, nitro, oxo, haloC₁₋₆ alkyl, polyhaloC₁₋₆ alkyl, haloC₁₋₆ alkoxy, polyhaloC₁₋₆ alkoxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkoxyC₁₋₆ alkyl, C₃₋₇ cycloalkylC₁₋₆ alkoxy, C₁₋₆ alkanoyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyloxy, C₁₋₆ alkylsulfonylC₁₋₆ alkyl, C₁₋₆ alkylsulfonamidoC₁₋₆ alkyl, C₁₋₆ alkylamidoC₁₋₆ alkyl, arylsulfonyl, arylsulfonyloxy, aryloxy, arylsulfonamido, arylcarboxamido, aroyl, or a group NR¹⁵R¹⁶, -CONR¹⁵R¹⁶, -NR¹⁵COR¹⁶, -NR¹⁵SO₂R¹⁶ or -SO₂NR¹⁵R¹⁶, wherein R¹⁵ and R¹⁶ independently represent hydrogen or C₁₋₆ alkyl or together form a heterocyclic ring; X represents a bond, O, CO, SO₂, OCH₂ or CH₂O; each R² independently represents C₁₋₄ alkyl; R³ represents C₃₋₈ alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, C₃₋₆ cycloalkyl, C₅₋₆ cycloalkenyl or -C₁₋₄alkyl-C₃₋₆ cycloalkyl; wherein said C₃₋₆ cycloalkyl groups of R³ may be optionally substituted by one or more (eg. 1, 2 or 3) substituents which may be the same or different, and which are selected from the group consisting of halogen, C₁₋₄ alkyl or trifluoromethyl groups; n independently represents 0, 1 or 2; p is 1 or 2; or a pharmaceutically acceptable salt thereof.

- Specific compounds of formula (I) which may be mentioned are those wherein R¹ represents pyridyl or pyrimidyl optionally substituted by one or two hydrogen, amino, halogen, cyano, C₁₋₆ alkyl or C₁₋₆ alkoxy groups and R³ represents C₃₋₆ alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, C₃₋₆ cycloalkyl or C₅₋₆ cycloalkenyl.

- Alkyl groups, whether alone or as part of another group, may be straight chain or branched and the groups alkoxy and alkanoyl shall be interpreted similarly. The term 'halogen' is used herein to describe, unless otherwise stated, a group selected from fluorine, chlorine, bromine or iodine and the term 'polyhalo' is used herein to refer to a moiety containing more than one (eg. 2-5) of said halogen atoms.

- The term "aryl" includes single and fused rings wherein at least one ring is aromatic, for example, phenyl, naphthyl and tetrahydronaphthalenyl.

- The term "heterocyclyl" is intended to mean a 4-7 membered monocyclic saturated or partially unsaturated aliphatic ring or a 4-7 membered saturated or partially unsaturated aliphatic ring fused to a benzene ring containing 1 to 3 heteroatoms selected from oxygen, nitrogen or sulphur. Suitable examples of such monocyclic rings include pyrrolidinyl, azetidiny, piperidinyl, piperazinyl, morpholinyl, tetrahydrofuranly,

tetrahydropyranyl, diazepanyl and azepanyl. Suitable examples of benzofused heterocyclic rings include indoliny, isoindoliny, 2,3,4,5-tetrahydro-1*H*-3-benzazepine or tetrahydroisoquinoliny.

- 5 The term "heteroaryl" is intended to mean a 5-6 membered monocyclic aromatic or a fused 8-10 membered bicyclic aromatic ring containing 1 to 3 heteroatoms selected from oxygen, nitrogen and sulphur. Suitable examples of such monocyclic aromatic rings include thienyl, furyl, pyrrolyl, triazolyl, imidazolyl, oxazolyl, thiazolyl, oxadiazolyl, isothiazolyl, isoxazolyl, thiadiazolyl, pyrazolyl, pyrimidyl, pyridazinyl, pyrazinyl and
10 pyridyl. Suitable examples of such fused aromatic rings include benzofused aromatic rings such as quinoliny, isoquinoliny, quinazoliny, quinoxaliny, cinnoliny, naphthyridiny, indolyl, indazolyl, pyrrolopyridiny, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzoxadiazolyl, benzothiadiazolyl and the like.

- 15 Preferably, R¹ represents
aryl (eg. phenyl) optionally substituted by one or more (eg. 1, 2 or 3) cyano or C₁₋₆ alkoxy carbonyl (eg. ethoxy carbonyl) groups;
heteroaryl (eg. pyridyl or pyrimidyl) optionally substituted by one or more (eg. 1, 2
20 or 3) cyano, halogen (eg. bromine), polyhaloC₁₋₆ alkyl (eg. CF₃) or C₁₋₆ alkoxy carbonyl (eg. methoxy carbonyl) groups;
-heteroaryl-X-heteroaryl (eg. pyrimidyl-pyridyl); or
-heteroaryl-X-heterocyclyl (eg. pyrimidyl-morpholiny).

- 25 More preferably, R¹ represents
aryl (eg. phenyl) optionally substituted by one or more (eg. 1, 2 or 3) cyano or C₁₋₆ alkoxy carbonyl (eg. ethoxy carbonyl) groups;
heteroaryl (eg. pyridyl or pyrimidyl) optionally substituted by one or more (eg. 1, 2
or 3) cyano, halogen (eg. bromine), polyhaloC₁₋₆ alkyl (eg. CF₃) or C₁₋₆ alkoxy carbonyl
30 (eg. methoxy carbonyl) groups.

Preferably X represents a bond.

Preferably, n represents 0.

35

Preferably, p represents 1.

Preferably, R³ represents C₃₋₈ alkyl (eg. isopropyl).


- 40 Preferred compounds according to the invention include examples E1-E8 as shown below, or a pharmaceutically acceptable salt thereof.

Compounds of formula (I) may form acid addition salts with acids, such as conventional pharmaceutically acceptable acids, for example maleic, hydrochloric, hydrobromic, phosphoric, acetic, fumaric, salicylic, sulphate, citric, lactic, mandelic, tartaric and methanesulphonic.

Certain compounds of formula (I) are capable of existing in stereoisomeric forms. It will be understood that the invention encompasses all geometric and optical isomers of these compounds and the mixtures thereof including racemates. Tautomers also form an aspect of the invention.

The present invention also provides a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which process comprises:


(a) reacting a compound of formula (II)



(II)


or an optionally activated or protected derivative thereof, wherein R^2 , n and p are as defined above and R^{3a} is as defined for R^3 above or a group convertible to R^3 , with a compound of formula R^1-L^1 , wherein R^1 is as defined above and L^1 represents a suitable leaving group, such as a halogen atom (eg. fluorine or chlorine) followed by a deprotection reaction as necessary; or

(b) reacting a compound of formula (III)



(III)

wherein R^1 is defined above and L^2 represents OH or a suitable leaving group, such as a halogen atom (eg. chlorine), with a compound of formula (IV)



(IV)

wherein R^2 , n and p are as defined above R^{3a} is as defined for R^3 above or a group convertible to R^3 ; or

5

(c) deprotecting a compound of formula (I) or converting groups which are protected; and optionally thereafter

(d) interconversion to other compounds of formula (I).

10

Process (a) typically comprises the use of a suitable base, such as potassium carbonate in a suitable solvent such as dimethylsulfoxide or N,N-dimethylformamide at elevated temperature.

15

Process (b) typically comprises activation of the compound of formula (III) wherein L^2 represents OH with a coupling reagent such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) in the presence of 1-hydroxybenzotriazole (HOBT) in a suitable solvent such as dichloromethane followed by reaction with the compound of formula (IV).

20

Process (b) may also involve halogenation of the compound of formula (III) with a suitable halogenating agent (eg. thionyl chloride) followed by reaction with the compound of formula (IV) in the presence of a suitable base such as triethylamine or a solid supported base such as diethylaminomethylpolystyrene in a suitable solvent such as dichloromethane

25

In process (c), examples of protecting groups and the means for their removal can be found in T. W. Greene 'Protective Groups in Organic Synthesis' (J. Wiley and Sons, 1991). Suitable amine protecting groups include sulphonyl (e.g. tosyl), acyl (e.g. acetyl),

30

2',2',2'-trichloroethoxycarbonyl, benzyloxycarbonyl or t-butoxycarbonyl) and arylalkyl (e.g. benzyl), which may be removed by hydrolysis (e.g. using an acid such as hydrochloric acid) or reductively (e.g. hydrogenolysis of a benzyl group or reductive removal of a 2',2',2'-trichloroethoxycarbonyl group using zinc in acetic acid) as appropriate. Other suitable amine protecting groups include trifluoroacetyl ($-COCF_3$)


35

which may be removed by base catalysed hydrolysis or a solid phase resin bound benzyl group, such as a Merrifield resin bound 2,6-dimethoxybenzyl group (Ellman linker), which may be removed by acid catalysed hydrolysis, for example with trifluoroacetic acid.

Process (d) may be performed using conventional interconversion procedures such as epimerisation, oxidation, reduction, alkylation, nucleophilic or electrophilic aromatic substitution, ester hydrolysis or amide bond formation.

5

Compounds of formula (II) may be prepared in accordance with the following procedure:



10 wherein R^2 , n and p are as defined above, R^{3a} is as defined for R^3 above or a group convertible to R^3 , L^3 represents OH or a suitable leaving group such as a halogen atom (eg. chlorine), and P^1 represents a suitable protecting group such as t-butoxycarbonyl.


15 When L^3 represents OH, step (i) typically comprises the use of suitable coupling conditions eg 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) in the presence of 1-hydroxybenzotriazole (HOBT) or 1-hydroxy-7-azabenzotriazole (HOAT).

When L^3 represents a suitable leaving group such as a halogen atom (eg. chlorine), step (i) typically comprises the use of a suitable base such as triethylamine in a suitable solvent such as dichloromethane.

20

Step (ii) typically comprises a suitable deprotection reaction using standard conditions such as those described above for process (c). Where P^1 is a tert butoxycarbonyl group this may involve a suitable acid such as HCl or trifluoroacetic acid

Compounds of formula (III) wherein L^2 represents OH, may be prepared in accordance with the following procedure:



5

Step (i) is typically carried out in a suitable solvent such as N,N-dimethylformamide in the presence of a base such as potassium carbonate.

- 10 Step (ii) typically comprises a suitable deprotection reaction using standard conditions such as those described above for process (c). Where P^2 is an alkoxy group such as ethoxy this may involve a suitable acid such as HCl or a base such as sodium hydroxide.

- 15 Compounds of formula (III) wherein L^2 represents a suitable leaving group, such as a halogen atom (eg. chlorine) may be prepared by treating a compound of formula (III)^a with thionyl chloride or oxalyl chloride.

Compounds of formula (IV) and (V) are either known in the literature or can be prepared by analogous methods.

20

- Compounds of formula (I) and their pharmaceutically acceptable salts have affinity for the histamine H₃ receptor and are believed to be of potential use in the treatment of neurological diseases including Alzheimer's disease, dementia, age-related memory dysfunction, mild cognitive impairment, cognitive deficit, epilepsy, neuropathic pain, inflammatory pain, Parkinson's disease, multiple sclerosis, stroke and sleep disorders including narcolepsy; psychiatric disorders including schizophrenia (particularly cognitive deficit of schizophrenia), attention deficit hyperactivity disorder, depression and addiction; and other diseases including obesity, asthma, allergic rhinitis, nasal congestion, chronic obstructive pulmonary disease and gastro-intestinal disorders.

30

Thus the invention also provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use as a therapeutic substance in the treatment or

prophylaxis of the above disorders, in particular neurodegenerative disorders including Alzheimer's disease.

5 The invention further provides a method of treatment or prophylaxis of the above disorders, in mammals including humans, which comprises administering to the sufferer a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

10 In another aspect, the invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the treatment of the above disorders.

15 When used in therapy, the compounds of formula (I) are usually formulated in a standard pharmaceutical composition. Such compositions can be prepared using standard procedures.

20 Thus, the present invention further provides a pharmaceutical composition for use in the treatment of the above disorders which comprises the compound of formula (I) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

25 The present invention further provides a pharmaceutical composition which comprises the compound of formula (I) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

30 Compounds of formula (I) may be used in combination with other therapeutic agents, for example histamine H1 antagonists or medicaments claimed to be useful as either disease modifying or symptomatic treatments of Alzheimer's disease. Suitable examples of such other therapeutic agents may be agents known to modify cholinergic transmission such as 5-HT₆ antagonists, M1 muscarinic agonists, M2 muscarinic antagonists or acetylcholinesterase inhibitors. When the compounds are used in combination with other therapeutic agents, the compounds may be administered either sequentially or simultaneously by any convenient route.

35 The invention thus provides, in a further aspect, a combination comprising a compound of formula (I) or a pharmaceutically acceptable derivative thereof together with a further therapeutic agent or agents.

40 The combinations referred to above may conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical formulations comprising a combination as defined above together with a pharmaceutically acceptable carrier or excipient comprise a further aspect of the invention. The individual components of such

combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations.

5 When a compound of formula (I) or a pharmaceutically acceptable derivative thereof is used in combination with a second therapeutic agent active against the same disease state the dose of each compound may differ from that when the compound is used alone. Appropriate doses will be readily appreciated by those skilled in the art.

10 A pharmaceutical composition of the invention, which may be prepared by admixture, suitably at ambient temperature and atmospheric pressure, is usually adapted for oral, parenteral or rectal administration and, as such, may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable or infusible solutions or suspensions or suppositories. Orally administrable compositions are generally preferred.

15 Tablets and capsules for oral administration may be in unit dose form, and may contain conventional excipients, such as binding agents, fillers, tableting lubricants, disintegrants and acceptable wetting agents. The tablets may be coated according to methods well known in normal pharmaceutical practice.

20 Oral liquid preparations may be in the form of, for example, aqueous or oily suspension, solutions, emulsions, syrups or elixirs, or may be in the form of a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents,
25 non-aqueous vehicles (which may include edible oils), preservatives, and, if desired, conventional flavourings or colorants.

For parenteral administration, fluid unit dosage forms are prepared utilising a compound of the invention or pharmaceutically acceptable salt thereof and a sterile vehicle. The
30 compound, depending on the vehicle and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions, the compound can be dissolved for injection and filter sterilised before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are dissolved in the vehicle. To enhance the stability, the composition can be
35 frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is suspended in the vehicle instead of being dissolved, and sterilisation cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before suspension in a sterile vehicle. Advantageously, a surfactant or wetting
40 agent is included in the composition to facilitate uniform distribution of the compound.

- The composition may contain from 0.1% to 99% by weight, preferably from 10 to 60% by weight, of the active material, depending on the method of administration. The dose of the compound used in the treatment of the aforementioned disorders will vary in the usual way with the seriousness of the disorders, the weight of the sufferer, and other similar factors. However, as a general guide suitable unit doses may be 0.05 to 1000 mg, more suitably 1.0 to 200 mg, and such unit doses may be administered more than once a day, for example two or three a day. Such therapy may extend for a number of weeks or months.
- The following Descriptions and Examples illustrate the preparation of compounds of the invention.


Description 1

4-[(4-Isopropyl-piperazin-1-yl)carbonyl]-piperidine (D1)

- 1-*tert*-Butoxycarbonyl-piperidine-4-carboxylate (5g) in DMF (60ml) was treated with EDC (5.5g) followed by HOAT (0.1g). After 5min, N-isopropylpiperazine (2.8g) in DMF (5ml) was added and the reaction was left stirring for 18h. The reaction was then diluted with EtOAc (150ml) and washed with saturated brine/sodium hydrogen carbonate (1:1, 200ml) followed by brine (3x200ml). The EtOAc layer was evaporated to near dryness and the residue treated with TFA/water (40ml, 95:5) for 5h prior to evaporation and re-evaporation from toluene (3x60ml). The residue was taken up into a minimum volume of EtOAc and treated with HCl (30ml, 2N solution in diethyl ether) for 1h. The resulting dihydrochloride salt was filtered off and washed with diethyl ether before drying under vacuum (4.8g). Dissolving the dihydrochloride in water and basifying with potassium carbonate, followed by extraction with EtOAc and evaporation gave the title compound (D1) as the free base (3.5g). ¹H NMR δ [CD₃OD]; 1.073 (6H, d, J= 6.4Hz), 1.69-1.75 (4H, m), 2.5-2.58 (4H, m), 2.69-2.82 (3H, m), 2.85-2.94 (1H, m), 3.13-3.22 (2H, m) and 3.54-3.65 (4H, m).


Example 1

N-(5-Cyano-pyridin-2-yl)-4-[4-(isopropyl)piperazin-1-yl]carbonyl]-piperidine hydrochloride (E1)



- To 4-[4-(isopropyl)piperazin-1-ylcarbonyl]-piperidine (0.239g) (D1) and 2-chloro-5-cyano-pyridine (0.138g), dissolved in DMSO (5ml), was added potassium carbonate (0.14g). The reaction was heated to 80°C for 4h before cooling and dilution with saturated sodium hydrogen carbonate (50ml) and EtOAc (80ml). The EtOAc layer was washed further with brine (3x80ml) and then extracted with 1N HCl. The aqueous HCl extract was basified and back-extracted with EtOAc which was concentrated under vacuum. A solution of 2N HCl in diethyl ether (1ml) was then added and the precipitate filtered and washed with


diethyl ether. Crystallisation from methanol afforded the title compound (E1) (0.15g). ¹H NMR δ (DMSO-d₆): 1.29 (6H, d, J= 6.4Hz), 1.4-1.6 (2H, m), 1.7-1.85 (2H, m), 2.8-3.27 (6H, m), 3.32-3.52 (3H, m), 3.69 (1H, m), 4.24 (1H, brd, J=13Hz), 4.35-4.55 (3H, m), 6.96 (1H, d, J=9.1Hz), 7.84 (1H, dd, J=9.1Hz and 1.5Hz), 8.47 (1H, d, J=1.5Hz) and 11.36 (1H, brs). MS electrospray; (+ve ion) 342 (MH⁺).

Example 2**N-(4-Cyanophenyl)-4-[4-(isopropyl)piperazin-1-yl]carbonyl]-piperidine hydrochloride (E2)**

The title compound (E2) was prepared from 4-fluorobenzonitrile and 4-[4-(isopropyl)piperazin-1-ylcarbonyl]-piperidine (D1) according to the procedure described in Example 1, except that the reaction was carried out at 120°C for 8h. ¹H NMR δ (DMSO-d₆): 1.29 (6H, d, J= 6.5 Hz), 1.4-1.8 (4H, m), 2.8-3.1 (5H, m), 3.11-3.22 (1H, m), 3.31-3.5 (3H, m), 3.62-3.77 (1H, m), 3.89-4 (2H, m), 4.2 (1H, brd, J=13.5Hz), 7.03 (2H, d, J=9Hz), 7.56 (2H, d, J=9Hz) and 11.43 (1H, brs). MS electrospray; (+ve ion) 341 (MH⁺).

Examples 3-5 (E3-E5)


Examples 3 - 5 were prepared from 4-[4-(isopropyl)piperazin-1-yl]carbonyl]-piperidine (D1) and the appropriate heteroaryl chloride using the procedure described in Example 1 and displayed ¹H NMR and mass spectral data that were consistent with structure.



Example No	R	Heteroaryl Chloride	Mass Spectrum (ES ⁺)
E3			[MH] ⁺ 396/398
E4			[MH] ⁺ 443
E5			[MH] ⁺ 342

Examples 6-8 (E6-E8)

Examples 6 - 8 were prepared from 4-[4-(isopropyl)piperazin-1-ylcarbonyl]-piperidine (D1) and the appropriate aryl fluoride using the procedure described in Example 2 and displayed ^1H NMR and mass spectral data that were consistent with structure.



Example No	R	Aryl Fluoride	Mass Spectrum (ES^+)
E6			$[\text{MH}]^+$ 341
E7			$[\text{MH}]^+$ 341
E8			$[\text{MH}]^+$ 388

Abbreviations

DMSO	dimethylsulfoxide
DMF	N,N-dimethylformamide
EDC	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
EtOAc	ethyl acetate
HOAT	1-hydroxy-7-azabenzotriazole
h	hour
min	minutes
rt	room temperature
TFA	trifluoroacetic acid

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

Biological Data

A membrane preparation containing histamine H3 receptors may be prepared in accordance with the following procedures:

(i) Generation of histamine H3 cell line

DNA encoding the human histamine H3 gene was cloned into a holding vector, pCDNA3.1 TOPO (Invitrogen) and its cDNA was isolated from this vector by restriction digestion of plasmid DNA with the enzymes BamH1 and Not-1 and ligated into the inducible expression vector pGene (Invitrogen) digested with the same enzymes. The GeneSwitch™ system (a system where in transgene expression is switched off in the

absence of an inducer and switched on in the presence of an inducer) was performed as described in US Patent nos: 5,364,791; 5,874,534; and 5,935,934. Ligated DNA was transformed into competent DH5 α *E. coli* host bacterial cells and plated onto Luria Broth (LB) agar containing Zeocin™ (an antibiotic which allows the selection of cells expressing the *sh ble* gene which is present on pGene and pSwitch) at 50 μ g ml⁻¹. Colonies containing the re-ligated plasmid were identified by restriction analysis. DNA for transfection into mammalian cells was prepared from 250ml cultures of the host bacterium containing the pGeneH3 plasmid and isolated using a DNA preparation kit (Qiagen Midi-Prep) as per manufacturers guidelines (Qiagen).

CHO K1 cells previously transfected with the pSwitch regulatory plasmid (InVitrogen) were seeded at 2x10⁶ cells per T75 flask in Complete Medium, containing Hams F12 (GIBCOBRL, Life Technologies) medium supplemented with 10% v/v dialysed foetal bovine serum, L-glutamine, and hygromycin (100 μ g ml⁻¹), 24 hours prior to use. Plasmid DNA was transfected into the cells using Lipofectamine plus according to the manufacturers guidelines (InVitrogen). 48 hours post transfection cells were placed into complete medium supplemented with 500 μ g ml⁻¹ Zeocin™.

10-14 days post selection 10nM Mifepristone (InVitrogen), was added to the culture medium to induce the expression of the receptor. 18 hours post induction cells were detached from the flask using ethylenediamine tetra-acetic acid (EDTA; 1:5000; InVitrogen), following several washes with phosphate buffered saline pH 7.4 and resuspended in Sorting Medium containing Minimum Essential Medium (MEM), without phenol red, and supplemented with Earles salts and 3% Foetal Clone II (Hyclone). Approximately 1x 10⁷ cells were examined for receptor expression by staining with a rabbit polyclonal antibody, 4a, raised against the N-terminal domain of the histamine H3 receptor, incubated on ice for 60 minutes, followed by two washes in sorting medium. Receptor bound antibody was detected by incubation of the cells for 60 minutes on ice with a goat anti rabbit antibody, conjugated with Alexa 488 fluorescence marker (Molecular Probes). Following two further washes with Sorting Medium, cells were filtered through a 50 μ m Filcon™ (BD Biosciences) and then analysed on a FACS Vantage SE Flow Cytometer fitted with an Automatic Cell Deposition Unit. Control cells were non-induced cells treated in a similar manner. Positively stained cells were sorted as single cells into 96-well plates, containing Complete Medium containing 500 μ g ml⁻¹ Zeocin™ and allowed to expand before reanalysis for receptor expression via antibody and ligand binding studies. One clone, 3H3, was selected for membrane preparation.

(ii) Membrane preparation from cultured cells

All steps of the protocol are carried out at 4°C and with pre-cooled reagents. The cell pellet is resuspended in 10 volumes of buffer A2 containing 50mM N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES) (pH 7.40) supplemented with 10e-4M leupeptin (acetyl-leucyl-leucyl-arginal; Sigma L2884), 25 μ g/ml bacitracin (Sigma B0125), 1mM ethylenediamine tetra-acetic acid (EDTA), 1mM phenylmethylsulfonyl fluoride (PMSF) and 2x10e-6M pepstain A (Sigma). The cells are then homogenised by

- 2 x 15 second bursts in a 1 litre glass Waring blender, followed by centrifugation at 500g for 20 minutes. The supernatant is then spun at 48,000g for 30 minutes. The pellet is resuspended in 4 volumes of buffer A2 by vortexing for 5 seconds, followed by homogenisation in a Dounce homogeniser (10-15 strokes). At this point the preparation is aliquoted into polypropylene tubes and stored at -70°C.

Compounds of the invention may be tested for *in vitro* biological activity in accordance with the following assays:

10 (I) **Histamine H3 binding assay**

For each compound being assayed, in a white walled clear bottom 96 well plate, is added:-

- (a) 10µl of test compound (or 10µl of iodophenpropit (a known histamine H3 antagonist) at a final concentration of 10mM) diluted to the required concentration in 10% DMSO;
 - (b) 10µl ¹²⁵I 4-[3-(4-iodophenylmethoxy)propyl]-1H-imidazolium (iodoproxyfan) (Amersham; 1.85MBq/µl or 50µCi/ml; Specific Activity ~2000Ci/mmol) diluted to 200pM in assay buffer (50mM Tris(hydroxymethyl)aminomethane buffer (TRIS) pH 7.4, 0.5mM ethylenediamine tetra-acetic acid (EDTA)) to give 20pM final concentration; and
 - (c) 80µl bead/membrane mix prepared by suspending Scintillation Proximity Assay (SPA) bead type WGA-PVT at 100mg/ml in assay buffer followed by mixing with membrane (prepared in accordance with the methodology described above) and diluting in assay buffer to give a final volume of 80µl which contains 7.5µg protein and 0.25mg bead per well – mixture was pre-mixed at room temperature for 60 minutes on a roller.
- The plate is shaken for 5 minutes and then allowed to stand at room temperature for 3-4 hours prior to reading in a Wallac Microbeta counter on a 1 minute normalised tritium count protocol. Data was analysed using a 4-parameter logistic equation.

(II) **Histamine H3 functional antagonist assay**

- For each compound being assayed, in a white walled clear bottom 96 well plate, is added:-

- (a) 10µl of test compound (or 10µl of guanosine 5'- triphosphate (GTP) (Sigma) as non-specific binding control) diluted to required concentration in assay buffer (20mM N-2-Hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES) + 100mM NaCl + 10mM MgCl₂, pH7.4 NaOH);
- (b) 60µl bead/membrane/GDP mix prepared by suspending wheat germ agglutinin-polyvinyltoluene (WGA-PVT) scintillation proximity assay (SPA) beads at 100mg/ml in assay buffer followed by mixing with membrane (prepared in accordance with the methodology described above) and diluting in assay buffer to give a final volume of 60µl which contains 10µg protein and 0.5mg bead per well – mixture is pre-mixed at 4°C for 30 minutes on a roller and just prior to addition to the plate, 10µM final concentration of guanosine 5' diphosphate (GDP) (Sigma; diluted in assay buffer) is added;

The plate is incubated at room temperature to equilibrate antagonist with receptor/beads by shaking for 30 minutes followed by addition of:

- (c) 10 μ l histamine (Tocris) at a final concentration of 0.3 μ M; and
- (d) 20 μ l guanosine 5' [γ 35-S] thiotriphosphate, triethylamine salt (Amersham; radioactivity concentration = 37kBq/ μ l or 1mCi/ml; Specific Activity 1160Ci/mmol) diluted to 1.9nM in assay buffer to give 0.38nM final.


- 5
- 10
- The plate is then incubated on a shaker at room temperature for 30 minutes followed by centrifugation for 5 minutes at 1500 rpm. The plate is read between 3 and 6 hours after completion of centrifuge run in a Wallac Microbeta counter on a 1 minute normalised tritium count protocol. Data is analysed using a 4-parameter logistic equation. Basal activity used as minimum i.e. histamine not added to well.

Results

- 15
- The compounds of Examples E1-E8 were tested in the histamine H3 functional antagonist assay and exhibited pK_b values > 7.5. More particularly, the compounds of E1 E2, E4 and E8 exhibited pK_b values > 8.5.

CLAIMS

1. A compound of formula (I) or a pharmaceutically acceptable salt thereof:



(I)

wherein:

R¹ represents aryl, heteroaryl, -aryl-X-aryl, -aryl-X-heteroaryl, -aryl-X-heterocyclyl, -heteroaryl-X-heteroaryl, -heteroaryl-X-aryl or -heteroaryl-X-heterocyclyl;

wherein said aryl, heteroaryl and heterocyclyl groups of R¹ may be optionally substituted by one or more (eg. 1, 2 or 3) substituents which may be the same or different, and which are selected from the group consisting of halogen, hydroxy, cyano, nitro, oxo, haloC₁₋₆ alkyl, polyhaloC₁₋₆ alkyl, haloC₁₋₆ alkoxy, polyhaloC₁₋₆ alkoxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkoxyC₁₋₆ alkyl, C₃₋₇ cycloalkylC₁₋₆ alkoxy, C₁₋₆ alkanoyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyloxy, C₁₋₆ alkylsulfonylC₁₋₆ alkyl, C₁₋₆ alkylsulfonamidoC₁₋₆ alkyl, C₁₋₆ alkylamidoC₁₋₆ alkyl, arylsulfonyl, arylsulfonyloxy, aryloxy, arylsulfonamido, arylcarboxamido, aroyl, or a group NR¹⁵R¹⁶, -CONR¹⁵R¹⁶, -NR¹⁵COR¹⁶, -NR¹⁵SO₂R¹⁶ or -SO₂NR¹⁵R¹⁶, wherein R¹⁵ and R¹⁶ independently represent hydrogen or C₁₋₆ alkyl or together form a heterocyclic ring;

X represents a bond, O, CO, SO₂, OCH₂ or CH₂O;

each R² independently represents C₁₋₄ alkyl;

R³ represents C₃₋₈ alkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, C₃₋₆ cycloalkyl, C₅₋₆ cycloalkenyl or -C₁₋₄alkyl-C₃₋₆ cycloalkyl;

wherein said C₃₋₆ cycloalkyl groups of R³ may be optionally substituted by one or more (eg. 1, 2 or 3) substituents which may be the same or different, and which are selected from the group consisting of halogen, C₁₋₄ alkyl or trifluoromethyl groups;

n independently represents 0, 1 or 2;

p is 1 or 2;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 which is a compound of formula E1-E8 or a pharmaceutically acceptable salt thereof.

3. A compound according to claim 1 or claim 2 for use in therapy.

4. A compound according to claim 1 or claim 2 for use in the treatment of Alzheimer's disease.

PB60227P

5. A pharmaceutical composition which comprises a compound according to claim 1 or claim 2 and a pharmaceutically acceptable carrier or excipient.

5